arXiv:1304.6896v1 [math.CO] 25 Apr 2013

Some new light subgraphs in the 1-planar graphs

Tao Wang [a, b, *]
[a]Institute of Applied Mathematics
Henan University, Kaifeng, 475004, P. R. China
[b]College of Mathematics and Information Science
Henan University, Kaifeng, 475004, P. R. China

April 26, 2013

Abstract

A graph is 1-*planar* if it can be drawn on the plane such that every edge cross at most one other edge. A connected graph H is *strongly light* in a family of graphs $\mathfrak{G}$, if there exists a constant λ, such that every graph in $\mathfrak{G}$ contains a subgraph K isomorphic to H with $\deg_G(v) \leq \lambda$ for all $v \in V(K)$. In this paper, we present some strongly light subgraphs in the family of 1-planar graphs with minimum degree 7.

1 introduction

All graphs considered are finite, simple and undirected unless otherwise stated. Let G be a graph with vertex set V and edge set E. The *neighborhood* of a vertex v in a graph G, denoted by $N_G(v)$, is the set of all the vertices adjacent to the vertex v, i.e., $N_G(v) = \{u \in V(G) \mid uv \in E(G)\}$. The *degree* of a vertex v in G, denoted by $\deg_G(v)$, is the number of edges of G incident with v. We denote the minimum and maximum degrees of vertices of G by $\delta(G)$ and $\Delta(G)$, respectively. A vertex of degree k is called a k-vertex, and a vertex of degree at most or at least k is call a k^-- or k^+-vertex, respectively.

A graph is 1-*embeddable* in a surface S if it can be drawn on S such that every edge cross at most one other edge. In particular, a graph is 1-*toroidal* if it can be drawn on torus such that every edge cross at most one other edge; a graph is 1-*planar* if it can be drawn on the plane such that every edge cross at most one other edge. The concept of 1-planar graph was introduced by Ringel [8] in 1965, while he simultaneously color the vertices and faces of a plane graph such that any pair of adjacent/incident elements receive different colors. Ringel [8] proved that 1-planar graphs are 7-colorable, and conjectured that they are 6-colorable, this conjecture was proved to be true by Borodin [1, 2]. For the various coloring problem of 1-planar graphs, see [4, 10, 13].

A connected graph H is *strongly light* in a family of graphs $\mathfrak{G}$, if there exists a constant λ, such that every graph in $\mathfrak{G}$ contains a subgraph K isomorphic to H with $\deg_G(v) \leq \lambda$ for all $v \in V(K)$.

Fabrici and Madaras [5] studied the structure of 1-planar graphs, mainly on the light subgraphs of 1-planar graphs. It is shown that every 3-connected 1-planar graph contains an edge with both ends of degree at most 20, and this bound is best possible. For some other light subgraphs, we refer the reader to [3, 6, 9].

2 Light subgraphs

Let G be a graph having been drawn on a surface, if we treat all the crossing points as vertices, and then obtained an embedded graph $G^{\dagger}$, and call it *the associated graph of* G, call the vertices of G true vertices and the crossing points *crossing vertices*.

Proposition 1. If a graph G is light in a family of graphs $\mathfrak{G}$, then every connected subgraph of G is also light in $\mathfrak{G}$.

Theorem 2.1. There is no light connected graph in the family of 1-planar graphs.

Proof. Let G be an arbitrary connected 1-planar graph. There exists a drawing such that there are at least two true vertices w_1 and w_2 of G on the outer curve. Moreover, we can assume that $\deg_G(w_1) \geq 2$. Make n copies of this 1-diagram of G, and identify the corresponding vertex w_1, and identify the corresponding vertex w_2, the obtained

---

[*] `Corresponding author: wangtao@henu.edu.cn`

graph is 1-diagram of a 1-planar graph G_n (Note that we delete the multiple edge between u and v). In the graph $G_n - \{w_1, w_2\}$, every component has at most $|V(G)| - 2$ vertices, hence $G \nsubseteq G_n - \{w_1, w_2\}$. That is, every subgraph of G_n which is isomorphic to G must contains at least one vertex in $\{w_1, w_2\}$, this vertex has degree at least n in G_n, hence G is not a light graph in the family of 1-planar graphs. □

Zhang et al. have proved the following two results on light graphs.

Theorem 2.2 ([14])**.** Each 1-planar graph with minimum degree 7 contains a copy of K_4 with all vertices of degree at most 11.

Theorem 2.3 ([11])**.** Each 1-planar graph with minimum degree at least 7 contains a 4-cycle $C = [x_1x_2x_3x_4]$ of the type $(7, 10^-, 8^-, 10^-)$ with a chord x_1x_3.

We improve the above two results in some sense.

Theorem 2.4. If G is a 1-planar graph with minimum degree 7, then it contains a K_4 of the type $(7, 8^-, 8^-, 10^-)$.

Proof. Let G be a connected counterexample to the theorem and it has been optimal 1-embedded on the plane (that is, every face of its associated graph is homeomorphic to an open disc). Hence, the graph G contains no K_4 or every copy of K_4 is the type $(8^+, 8^+, 8^+, 8^+)$ or $(7, 9^+, 9^+, 9^+)$ or $(7, 8^-, 9^+, 9^+)$ or $(7, 8^-, 8^-, 11^+)$.

Let $K^\dagger$ be the associated graph of G. By Euler's formula, we have

$$\sum_{v \in V(K^\dagger)} (\deg_{K^\dagger}(v) - 6) + \sum_{f \in F(K^\dagger)} (2 \deg_{K^\dagger}(f) - 6) = -12. \tag{1}$$

We will use discharging method to complete the proof. The initial charge of every vertex v is $\deg_{K^\dagger}(v) - 6$, and the initial charge of every face f is $2 \deg_{K^\dagger}(f) - 6$. By (1), the sum of all the elements' charge is -12. We then transfer some charge from the big faces and some big vertices to crossing vertices, such that the final charge of every crossing vertex becomes nonnegative and the final charge of every big vertex and face remains nonnegative, and thus the sum of the final charge of vertices and faces is nonnegative, which leads to a contradiction.

The Discharging Rules:

(R1) Every 4^+-face donates its redundant charge equally to incident crossing vertices;

(R2) Every 7^+-vertex donates its redundant charge equally to incident false 3-faces.

(R3) After applying R1 and R2, every false 3-face donates its redundant charge to the incident crossing vertex.

From the discharging rules, the final charge of every face is nonnegative, and so is every 7^+-vertex. So it suffices to consider the final charge of crossing vertices in $K^\dagger$.

Note that every 7^+-vertex v is incident with at most $2\lfloor \deg(v)/2 \rfloor$ false 3-faces. More formally, every 7-vertex sends at least $1/6$ to each of its incident false 3-face; every 8-vertex sends at least $1/4$ to each of its incident false 3-face; every 9-vertex sends at least $3/8$ to each of its incident false 3-face; every 10-vertex sends at least $2/5$ to each of incident false 3-face; every 11^+-vertex sends at least $1/2$ to each of its incident false 3-face. Let v be an arbitrary crossing vertex in $K^\dagger$, notice that the four neighbors of v are 7^+-vertices.

If v is incident with at least two 4^+-faces, then its final charge is greater than $4 - 6 + 2 \times 1 = 0$. If v is incident with exactly one 4^+-face, then its final charge is at least $4 - 6 + 1 + 6 \times 1/6 = 0$.

If there is no crossing vertex which is incident with four 3-faces, then the sum of the final charge is nonnegative, which derive a contradiction. So we may assume that v is incident with four 3-faces. It is obvious that the four neighbors of v induce a K_4 in G. If this K_4 is of the type $(8^+, 8^+, 8^+, 8^+)$, then the final charge of v is at least $4-6+8\times 1/4 = 0$; if this K_4 is of the type $(7, 9^+, 9^+, 9^+)$, then the final charge of v is at least $4-6+2\times 1/6+6\times 3/8 > 0$; if this K_4 is of the type $(7, 8^-, 9^+, 9^+)$, then its final charge is $4 - 6 + 4 \times 1/6 + 4 \times 3/8 > 0$; if this K_4 is of the type $(7, 8^-, 8^-, 11^+)$, then the final charge of v is at least $4 - 6 + 6 \times 1/6 + 2 \times 1/2 = 0$.

Finally, all the faces and vertices have nonnegative charge, which leads to a contradiction. □

Theorem 2.5. If G is a 1-planar graph with minimum degree 7, then G contains a subgraph as illustrated in Fig (a). Moreover, every w_i is a 23^--vertex and at most one of them is a 12^+-vertex.

Proof. Let G be a connected 1-planar graph with minimum degree 7, and it has been optimal 1-embedded on the plane (that is, every face of its associated graph is homeomorphic to an open disc). Let $K^\dagger$ be the associated graph of G.



(a)

Fig. 1

By Euler's formula, we have

$$\sum_{v\in V(K^{\dagger})}(\deg_{K^{\dagger}}(v)-4)+\sum_{f\in F(K^{\dagger})}(\deg_{K^{\dagger}}(f)-4)=-8. \tag{2}$$

We will use discharging method to complete the proof. The initial charge of every vertex v is $\deg_{K^{\dagger}}(v)-4$, and the initial charge of every face f is $\deg_{K^{\dagger}}(f)-4$. By (3), the sum of all the elements' charge is -8. We then transfer some charge from 7^+-vertices to 3-faces, such that the final charge of every element except 7-vertex is nonnegative, hence there exists a 7-vertex such that its final charge is negative and the local structure is desired.

The Discharging Rules:

(R1) Every 7^+-vertex w sends $1/2$ to each incident false 3-face and sends $1/3$ to each incident true 3-face;

(R2) Let f be a face with a face angle w_1ww_2 and $\deg(w)=k\geq 8$.

- (a) If f is a 3-face with $\deg(w_1)=7$ and $\deg(w_2)\geq 8$, then w sends $\frac{k-4}{k}-\frac{1}{3}$ to w_1 through f;
- (b) If f is a 3-face with $\deg(w_1)=\deg(w_2)=7$, then each of w_1 and w_2 receives $\frac{k-4}{2k}-\frac{1}{6}$ from w through f;
- (c) If f is a false 3-face with crossing vertex w_1, then w_2 receives $\frac{k-4}{2k}-\frac{1}{4}$ from w through f, while w sends $\frac{k-4}{2k}-\frac{1}{4}$ through the crossing vertex w_1 to the other neighbor in G;
- (d) If f is a 4^+-face and w_1 is a true vertex, then w sends $\frac{k-4}{2k}-\frac{1}{4}$ to w_1;
- (e) If f is a 4^+-face and w_1 is a crossing vertex, then w sends $\frac{k-4}{2k}-\frac{1}{4}$ through the crossing vertex w_1 to the other neighbor in G.

From the discharging rules, the final charge of every face is nonnegative, and so is every 8^+-vertex. Hence, there exists a 7-vertex w_0 such that its final charge is negative.

If w_0 is incident with at least one 4^+-face, then its final charge is nonnegative. So we may assume that w_0 is incident with seven 3-faces. Notice that the number of incident false 3-faces are even. If w_0 is incident with at most four false 3-faces, then its final charge is at least $7-4-4\times 1/2-3\times 1/3=0$. Hence, the vertex w_0 must be incident with six false 3-faces and one true face, and then w_0 receives less than $1/3$ from other vertices and faces.

Let $w_1w_0w_2$ be the true 3-face. If both w_1 and w_2 are 8^+-vertices, then w_0 receives at least $1/2-1/3$ from each of w_1 and w_2, thus w_0 receives at least $1/3$, a contradiction. So at least one of w_1 and w_2 must be a 7-vertex, we may assume that w_1 is a 7-vertex, see Fig (a).

If w_0 has a neighbor of degree at least 24 in G, then w_0 receives at least $5/6-1/2=1/3$ from this neighbor, which derive a contradiction. Therefore, all the neighbors of w_0 have degree at most 23 in G.

If w_0 has at least two 12^+-neighbors, then w_0 will receive at least $2\times(\frac{12-4}{12}-\frac{1}{2})=\frac{1}{3}$, which leads to a contradiction. Hence, the vertex w_0 has at most one 12^+-neighbor. □

Corollary 1 (Hudák and Madaras [7])**.** Every 1-planar graph with minimum degree 7 contains an edge of type $(7,7)$.

Corollary 2. Every 1-planar graph with minimum degree 7 contains a copy of $K_{1,7}$ with the center of degree 7 and the other vertices of degree at most 23.

Theorem 2.6. If G is a 1-planar graph with minimum degree 7, then G contains at least one subgraph as illustrated in Fig (b)–(i).



(b)



(c)



(d)



(e)



(f)



(g)



(h)



(i)

Proof. Let G be a connected a 1-planar graph with minimum degree 7, and it has been optimal 1-embedded on the plane (that is, every face of its associated graph is homeomorphic to an open disc). Let $K^\dagger$ be the associated graph of G.

By Euler's formula, we have

$$\sum_{v\in V(K^\dagger)} (\deg_{K^\dagger}(v) - 4) + \sum_{f\in F(K^\dagger)} (\deg_{K^\dagger}(f) - 4) = -8. \tag{3}$$

We will use discharging method to complete the proof. The initial charge of every vertex v is $\deg_{K^\dagger}(v) - 4$, and the initial charge of every face f is $\deg_{K^\dagger}(f) - 4$. By (3), the sum of all the elements' charge is -8. We then transfer some charge from 7^+-vertices to 3-faces, such that the final charge of every element except 7-vertex is nonnegative, hence there exists a 7-vertex such that its final charge is negative and the local structure is desired.

The Discharging Rules:

(R1) Every 7^+-vertex w sends $1/2$ to each incident false 3-face and sends $1/3$ to each incident true 3-face;

(R2) Let f be a face with a face angle w_1ww_2 and $\deg(w) = k \geq 8$.

- (a) If f is a 3-face with $\deg(w_1) = 7$ and $\deg(w_2) \geq 8$, then w sends $\frac{k-4}{k} - \frac{1}{3}$ to w_1 through f;
- (b) If f is a 3-face with $\deg(w_1) = \deg(w_2) = 7$, then each of w_1 and w_2 receives $\frac{k-4}{2k} - \frac{1}{6}$ from w through f;
- (c) If f is a false 3-face with crossing vertex w_1, then w sends $\frac{k-4}{k} - \frac{1}{2}$ to w_2;
- (d) If f is a 4^+-face and w_1 is a true vertex, then w sends $\frac{k-4}{2k} - \frac{1}{4}$ to w_1.

From the discharging rules, the final charge of every face is nonnegative, and so is every 8^+-vertex. Hence, there exists a 7-vertex w_0 such that its final charge is negative.

If w_0 is incident with at least one 4^+-face, then its final charge is nonnegative. So we may assume that w_0 is incident with seven 3-faces. Notice that the number of incident false 3-faces are even. If w_0 is incident with at most four false 3-faces, then its final charge is at least $7 - 4 - 4 \times 1/2 - 3 \times 1/3 = 0$. Hence, the vertex w_0 must be incident with six false 3-faces and one true face, and then w_0 receives less than $1/3$ from other vertices and faces.

Let $w_1w_0w_2$ be the true 3-face. If both w_1 and w_2 are 8^+-vertices, then w_0 receives at least $1/2 - 1/3$ from each of w_1 and w_2, thus w_0 receives at least $1/3$, which derive a contradiction. So at least one of w_1 and w_2 must be a 7-vertex, we may assume that w_1 is a 7-vertex, see Fig (a).

Case 1 Both $\deg(w_4)$ and $\deg(w_6)$ belong to $\{7, 8\}$. Since the vertex w_0 receives less than $\frac{1}{3}$ from the vertex w_2, it follows that $\frac{\deg(w_2)-4}{\deg(w_2)} - \frac{1}{2} + (\frac{\deg(w_2)-4}{2\deg(w_2)} - \frac{1}{6}) < \frac{1}{3}$ and $\deg(w_2) < 12$, see Fig (b).

Case 2 Exactly one of $\deg(w_4)$ and $\deg(w_6)$ belongs to $\{7,8\}$. If w_2 is a 7-vertex and $\max\{\deg(w_4),\deg(w_6)\} \geq 12$, then w_0 will receive at least $2 \times (\frac{2}{3} - \frac{1}{2}) = \frac{1}{3}$, which is a contradiction. If w_2 is a 8-vertex and $\max\{\deg(w_4),\deg(w_6)\} \geq 11$, then w_0 will receive at least $2 \times (\frac{7}{11} - \frac{1}{2}) + \frac{1}{12} > \frac{1}{3}$, a contradiction. If w_2 is a 9-vertex and $\max\{\deg(w_4),\deg(w_6)\} \geq 10$, then w_0 receives at least $2\times(\frac{3}{5}-\frac{1}{2})+(\frac{5}{9}-\frac{1}{2})+(\frac{5}{18}-\frac{1}{6}) = \frac{11}{30} > \frac{1}{3}$, which also leads to a contradiction. If w_2 is a 10^+-vertex, then w_0 receives at least $2\times(\frac{5}{9}-\frac{1}{2})+(\frac{3}{5}-\frac{1}{2})+(\frac{1}{5}-\frac{1}{6}) > \frac{1}{3}$, a contradiction. So we have that w_2 is a 9^--vertex. If w_2 is a 7-vertex, then $\max\{\deg(w_4),\deg(w_6)\} \leq 11$, see Fig (c); if w_2 is a 8-vertex, then $\max\{\deg(w_4),\deg(w_6)\} \leq 10$, see Fig (d) and Fig (f); if w_2 is a 9-vertex, then $\max\{\deg(w_4),\deg(w_6)\} = 9$, see Fig (e) and Fig (g).

Case 3 Both $\deg(w_4)$ and $\deg(w_6)$ are at least 9. If w_2 is a 9^+-vertex, then the vertex w_0 will receive at least $(\frac{5}{18} - \frac{1}{6}) + 5 \times (\frac{5}{9} - \frac{1}{2}) > \frac{1}{3}$, a contradiction. It follows that w_2 is a 7- or 8-vertex. If $\min\{\deg(w_4),\deg(w_6)\} \geq 10$, then the vertex w_0 will receive at least $4 \times (\frac{3}{5} - \frac{1}{2}) = \frac{2}{5} > \frac{1}{3}$, a contradiction. Hence, we have that $\min\{\deg(w_4),\deg(w_6)\} = 9$. If w_2 is a 7-vertex and $\max\{\deg(w_4),\deg(w_6)\} \geq 11$, then the vertex w_0 will receive at least $2\times(\frac{7}{11}-\frac{1}{2})+2\times(\frac{5}{9}-\frac{1}{2}) > \frac{1}{3}$, a contradiction. If w_2 is a 8-vertex and $\max\{\deg(w_4),\deg(w_6)\} \geq 10$, then w_0 will receive at least $2 \times (\frac{3}{5} - \frac{1}{2}) + 2 \times (\frac{5}{9} - \frac{1}{2}) + \frac{1}{12} > \frac{1}{3}$, which is a contradiction. Thus, if w_2 is a 7-vertex, then G contains a subgraph as illustrated in Fig (h); if w_2 is a 8-vertex, then G contains a subgraph as illustrated in Fig (i). □

Corollary 3. If G is a 1-planar graph with minimum degree 7, then G contains a triangle of type $(7,7,9^-)$.

As an immediate consequence, the following corollary is an improvement of Theorem 2.3.

Corollary 4. If G is a 1-planar graph with minimum degree 7, then G contains a 4-cycle $C = [x_1x_2x_3x_4]$ of the type $(7,9^-,7,9^-)$ with a chord x_1x_3.

Corollary 5 ([12])**.** If G is a 1-planar graph with minimum degree 7, then G contains a copy of $K_1 \vee (K_1 \cup K_2)$ with all vertices of degree at most 9.

References

[1] O. V. Borodin, Solution of the Ringel problem on vertex-face coloring of planar graphs and coloring of 1-planar graphs, Metody Diskret. Analiz. (1984) (41) 12–26, 108.

[2] O. V. Borodin, A new proof of the 6 color theorem, J. Graph Theory 19 (1995) (4) 507–521.

[3] O. V. Borodin, I. G. Dmitriev and A. O. Ivanova, The height of a cycle of length 4 in 1-planar graphs with minimal degree 5 without triangles, Diskretn. Anal. Issled. Oper. 15 (2008) (1) 11–16.

[4] O. V. Borodin, A. V. Kostochka, A. Raspaud and E. Sopena, Acyclic colouring of 1-planar graphs, Discrete Appl. Math. 114 (2001) (1-3) 29–41.

[5] I. Fabrici and T. Madaras, The structure of 1-planar graphs, Discrete Math. 307 (2007) (7-8) 854–865.

[6] D. Hudák and T. Madaras, On local structure of 1-planar graphs of minimum degree 5 and girth 4, Discuss. Math. Graph Theory 29 (2009) (2) 385–400.

[7] D. Hudák and T. Madaras, On local properties of 1-planar graphs with high minimum degree, Ars Math. Contemp. 4 (2011) (2) 245–254.

[8] G. Ringel, Ein Sechsfarbenproblem auf der Kugel, Abh. Math. Sem. Univ. Hamburg 29 (1965) (1) 107–117.

[9] Y. Suzuki, Re-embeddings of maximum 1-planar graphs, SIAM J. Discrete Math. 24 (2010) (4) 1527–1540.

[10] X. Zhang and G. Liu, On edge colorings of 1-planar graphs without adjacent triangles, Inform. Process. Lett. 112 (2012) (4) 138–142.

[11] X. Zhang and G. Liu, On the lightness of chordal 4-cycle in 1-planar graphs with high minimum degree, Ars Math. Contemp. (2014).

[12] X. Zhang, G. Z. Liu and J. L. Wu, Light subgraphs in the family of 1-planar graphs with high minimum degree, Acta Math. Sin. (Engl. Ser.) 28 (2012) (6) 1155–1168.

[13] X. Zhang and J.-L. Wu, On edge colorings of 1-planar graphs, Inform. Process. Lett. 111 (2011) (3) 124–128.

[14] X. Zhang, J.-L. Wu and G. Liu, New upper bounds for the heights of some light subgraphs in 1-planar graphs with high minimum degree, Discrete Math. Theor. Comput. Sci. 13 (2011) (3) 9–16.